


- source optical coherence tomography. *Retina*. 2016;36:499–516.
13. Matsumoto H, Kishi S, Mukai R, Akiyama H. Remodeling of macular vortex veins in pachychoroid neovascularopathy. *Sci Rep*. 2019;9:14689.
 14. Hiroe T, Kishi S. Dilatation of asymmetric vortex vein in central serous chorioretinopathy. *Ophthalmol Retina*. 2018;2:152–161.
 15. Terao N, Imanaga N, Wakugawa S, et al. Short axial length is related to asymmetric vortex veins in central serous chorioretinopathy. *Ophthalmol Sci*. 2021;1:100071.
 16. Matsumoto H, Hoshino J, Mukai R, et al. Vortex vein anastomosis at the watershed in pachychoroid spectrum diseases. *Ophthalmol Retina*. 2020;4:938–945.
 17. Choudhry N, Duker JS, Freund KB, et al. Classification and guidelines for widefield imaging: recommendations from the international widefield imaging study group. *Ophthalmol Retina*. 2019;3:843–849.
 18. Hayreh SS. In vivo choroidal circulation and its watershed zones. *Eye (Lond)*. 1990;4:273–289.
 19. Jung JJ, Yu DJG, Ito K, et al. Quantitative assessment of asymmetric choroidal outflow in pachychoroid eyes on ultra-widefield indocyanine green angiography. *Invest Ophthalmol Vis Sci*. 2020;61:50.
 20. Jeong S, Kang W, Noh D, et al. Choroidal vascular alterations evaluated by ultra-widefield indocyanine green angiography in central serous chorioretinopathy. *Graefes Arch Clin Exp Ophthalmol*. 2022;260:1887–1898.
 21. Mori K, Gehlbach PL, Yoneya S, Shimizu K. Asymmetry of choroidal venous vascular patterns in the human eye. *Ophthalmology*. 2004;111:507–512.
 22. Cheung CMG, Lee WK, Koizumi H, et al. Pachychoroid disease. *Eye (Lond)*. 2019;33:14–33.
 23. Matsumoto H, Hoshino J, Arai Y, et al. Quantitative measures of vortex veins in the posterior pole in eyes with pachychoroid spectrum diseases. *Sci Rep*. 2020;10:19505.
 24. Mehta N, Chong J, Tsui E, et al. Presumed foveal bacillary layer detachment in a patient with toxoplasmosis chorioretinitis and pachychoroid disease. *Retin Cases Brief Rep*. 2021;15:391–398.
 25. Ramtohul P, Engelbert M, Malclès A, et al. Bacillary layer detachment: multimodal imaging and histologic evidence of a novel optical coherence tomography terminology: literature review and proposed theory. *Retina*. 2021;41:2193–2207.
 26. Lee K, Ra H, Lee JH, et al. Classification of pachychoroid on optical coherence tomographic en face images using deep convolutional neural networks. *Transl Vis Sci Technol*. 2021;10:28.

Pictures & Perspectives



Large Colloid Drusen

A 35-year-old man presented with bilateral large colloid drusen. Ultrawide-field fundus photography revealed several large colloid drusen in the macula and periphery of the retina (A). Late-phase fluorescein angiography revealed hyperfluorescent drusen of varying size (B). OCT revealed multiple dome-shaped retinal pigment epithelial elevations with variable internal reflectivity (C). The upper OCT was acquired 6 years ago, whereas the lower OCT was acquired at a recent visit. The choroidal slab of widefield en face swept-source OCT angiography revealed that well-demarcated drusen had a hyperreflective center surrounded by a hyporeflective ring (D). The choroidal vessels were not dilated. (Magnified version of Figure A-D is available online at www.opthalmologyretina.org/).

WENWEN CHEN, MD, PhD

QING CHANG, MD, PhD

Eye Institute and Department of Ophthalmology, Eye & ENT Hospital, Fudan University, Shanghai, China