



### Unusual Retinal Hemorrhages in a Highly Myopic Patient

A 46-year-old woman with high myopia (axial length >27 mm in both eyes) was found to have a tessellated fundus with aneurysmal lesions and intraretinal hemorrhages in the left macula during a routine ophthalmic examination (A). Infrared imaging further accentuated the lesions and fluorescein angiography confirmed saccular aneurysmal vessels with minimal dye leakage (B, C). OCT identified outer and inner layer retinoschisis and OCT angiography revealed flow through the aneurysms and dilated vessels within the schisis cavity (D). This uncommon disorder is postulated to be a result of mechanical disruption forces from vitreous traction anteriorly and progressive staphylomatous posterior elongation. (Magnified version of Figure A–D is available online at [www.ophtalmologyretina.org](http://www.ophtalmologyretina.org))

KAI-LAN CHANG, MD<sup>1</sup>

NAN-KAI WANG, MD, PhD<sup>2,3,4</sup>

SHAWN H. TSAI, MD, MS<sup>1,5</sup>

<sup>1</sup>Department of Ophthalmology, Mackay Memorial Hospital, Taipei, Taiwan; <sup>2</sup>College of Medicine, School of Medicine, Chang Gung University, Taoyuan, Taiwan; <sup>3</sup>Department of Ophthalmology, Chang Gung Memorial Hospital, Linkou Medical Center, Taoyuan, Taiwan; <sup>4</sup>Department of Ophthalmology, Vagelos College of Physicians and Surgeons, Columbia University Irving Medical Center, Columbia University, New York, New York; <sup>5</sup>Department of Optometry, Mackay Medical College, New Taipei City, Taiwan