


Spiderweb-Like Vitreous Changes in a Vasoproliferative Tumor

Search for articles by this author

Property	Value
Status	
Version	
Ad File	
Disable Ads Flag	
Environment	
Moat Init	
Moat Ready	
Contextual Ready	
Contextual URL	
Contextual Initial Segments	
Contextual Used Segments	
AdUnit	
SubAdUnit	
Custom Targeting	
Ad Events	
Invalid Ad Sizes	



- Tomoyuki Ishibashi, MD

Tomoyuki Ishibashi

Affiliations

Department of Ophthalmology, Osaka Rosai Hospital Clinical Research Center for Optical Sensory Organ Disability, Osaka, Japan

- Kazuyuki Emi, MD, PhD

Kazuyuki Emi

Affiliations

Department of Ophthalmology, Osaka Rosai Hospital Clinical Research Center for Optical Sensory Organ Disability, Osaka, Japan

[Search for articles by this author](#)

A 24-year-old man with a history of a vasoproliferative tumor in his right eye presented with right vision loss that had progressed over 5 years. His right eye best-corrected visual acuity had decreased to 20/320. A confocal light-emitting diode-based retinal imaging system clearly showed a 3-dimensional, spiderweb-like proliferative change in the vitreous body above the posterior pole (**A**) and abnormal retinal vessels in the peripheral retina (**B**). OCT showed significant retinal folds with overlying dense epiretinal membrane (**C**). Two years after the removal of the vitreous proliferation via vitrectomy, the retinal folds had resolved (**D**), and his best-corrected visual acuity recovered to 20/32. (Magnified version of Figure **A-D** is available online at www.opthalmologyretina.org).

Article info

Publication history

Published online: January 27, 2024

Identification

DOI: <https://doi.org/10.1016/j.oret.2023.12.011>

Copyright

© 2023 by the American Academy of Ophthalmology

ScienceDirect

[Access this article on ScienceDirect](#)

Figures

Related Articles

[View Abstract](#)

